

NUTRIENT MANAGEMENT GUIDELINES FOR STATE PROPERTY AND COMMERCIALY MANAGED TURFGRASS



University of Maryland
Turfgrass Technical Update
TT - 115 March 2003

Dr. Thomas R. Turner, Turfgrass Specialist
University of Maryland Department of Natural Resource Sciences & Landscape Architecture

Nutrient management laws passed by the Maryland Legislature in 1998 require that University of Maryland nutrient management guidelines be followed on state property and commercially managed turfgrass sites. These laws impacting turfgrass are part of an overall effort to regulate the agricultural industry. However, major differences exist between the goals and practices of general agriculture and turfgrass management. The following information is intended to serve as a nutrient management guideline for the maintenance in an efficient, effective, and environmentally sound manner of state property and commercially maintained turfgrass sites.

Water Quality

Properly managed turfgrass has been shown to be an environmental asset. Turfgrass has significant cooling effects during the summer and traps much of the dust and dirt that is released each year into the atmosphere. Turfgrass absorbs carbon dioxide, ozone, sulfur dioxide, and other gases while releasing oxygen.

Water runoff is greatly reduced and water infiltration increased compared to most other agriculture and plant systems. Once turfgrass is established, soil loss from erosion is negligible. Runoff from established turf has compared favorably to forested land. Also, turfgrass is an efficient organic matter producing system. Thus, little nitrogen (N) or phosphorus (P) is lost from turfgrass sites if sound nutrient management practices are followed.

Onsite monitoring and numerous research studies have shown that nutrient loss from turfgrass sites is very small. However, research has also shown that certain types of improper N applications on specific types of sites can result in excessive nitrate (NO_3) leaching. This problem is very specific and has occurred as follows:

1. Very high rates of N were applied using soluble $\text{NO}_3\text{-N}$ containing fertilizers (e.g. ammonium nitrate [NH_4NO_3]),
2. The fertilizer was applied to dormant turf, such as bermudagrass during the winter,
3. Soils were predominantly sand, and
4. The sites had high water tables.

A sound turfgrass nutrient management plan considers this set of conditions that can cause a potential problem.

Nitrogen

Proper nitrogen fertilization is essential in maintaining quality turf that is resistant to pest problems, tolerant of stresses, and able to recover from damage. Current N recommendations are based on extensive research and are dependent on a variety of factors such as turfgrass species and cultivars, age of turf, soil type, management practices being used (irrigation, clipping removal, pest control programs), weather conditions, use of area, length of growing season, and the need for recovery from pest damage, adverse environmental conditions, and traffic.

The nutrient management specialist must take into account all these factors in devising an appropriate N fertilization program. The program may vary from year to year as these conditions change.

KEY POINTS

Take a soil test sometime during the first year of maintenance.

Take a soil test once every three years after the initial soil test.

Don't apply more than 1 lb. of soluble N per 1000 ft^2 in any one application.

Adjust annual nitrogen rates for turf species, age of turf, and use of turf.

Three major factors must be considered in developing an N application plan: 1. What types of N should be applied, 2. How much N must be applied during the year, and 3. When should N be applied?

Inadequate attention to each of these factors increases the potential for thin turf that is more prone to pest and stress problems. Thin turf also results in sites which are more prone to soil erosion. Additionally, the potential for leaching and/or runoff of N increases if guidelines are not followed.

Sources of Nitrogen

A wide range of N-containing fertilizers are available to the turfgrass manager. These fertilizers generally fall into one of two broad categories: 1) fertilizers that contain only soluble, quickly available N, or 2) fertilizers that contain some N in a slowly available form which is not immediately available for plant use.

Quickly available N-fertilizers contain $\text{NO}_3\text{-N}$ and/or $\text{NH}_4\text{-N}$, which are soluble and readily available for uptake by turfgrass plants. Turfgrass uptake may occur within a few days with $\text{NO}_3\text{-N}$ fertilizer. Nitrogen uptake may begin within 7 – 10 days with $\text{NH}_4\text{-N}$ fertilizers, as $\text{NH}_4\text{-N}$ is converted to $\text{NO}_3\text{-N}$ in the soil. Nitrogen uptake by turfgrass roots is predominately in the NO_3 form.

Leaching and runoff potential is much higher for $\text{NO}_3\text{-N}$. Thus, where conditions exist that are conducive to leaching or runoff, fertilizers that contain significant amounts of $\text{NO}_3\text{-N}$ should not be used. These conditions include sandy sites (sands and loamy sands) with high water tables when turf is not actively growing, and sites that are highly sloped. Fertilizers high in $\text{NO}_3\text{-N}$ include NH_4NO_3 , potassium nitrate, and calcium nitrate. Fertilizers that contain predominantly $\text{NO}_3\text{-N}$ should only be used on sites not prone to runoff or leaching, where very rapid response is essential, and on turf that is actively growing.

Soluble N fertilizers that contain $\text{NH}_4\text{-N}$ include urea ($\text{NH}_2\text{-CO-NH}_2$), ammonium sulfate (NH_4SO_4), and ammonium chloride (NH_4Cl). These fertilizers can produce excellent quality turf without leaching or runoff problems if used properly. Most important is limiting these fertilizers to 1.0 lb. N/1000 ft^2 per application. Higher rates per application can result in excessive growth of turf and can increase the potential for N loss on some sites.

Slow release fertilizers contain significant amounts of N that are not immediately available for plant uptake. Examples of fertilizer sources that contain various amounts of slow release N includes sulfur coated ureas, polymer coated ureas, methylene ureas, ureaformaldehydes, IBDU, natural organics, and various types of sludge. The N in all slow release fertilizers used for turfgrass maintenance is $\text{NH}_4\text{-N}$ based. Slow release fertilizers, while varying considerably in individual characteristics and release patterns, typically provide more even turfgrass response, provide N over a longer period of time, and are less prone to N leaching and runoff as compared to soluble N fertilizers. Their use should be considered on sites that are prone to leaching or runoff, and when an N application needs to be made to turfgrass during non-optimum growing conditions.

Rates of Nitrogen

When discussing rates of N fertilization, there are two primary issues: how much N can be applied in any one application, and how much total N can be applied annually. The maximum amount of N that should be applied in one application is primarily dependent on the amount of soluble N in the fertilizer. No more than 1.0 lb soluble N/1000 ft^2 should be made in any single application. For a fertilizer that contains 50% of its N in a soluble form and 50% in a slow release (water insoluble) form, no more than 2.0 lbs. total N/1000 ft^2 should thus be applied in any single application (which would result in 1.0 lb. soluble N/1000 ft^2 being applied).

Because some fertilizers contain most of their N in a slow release form, the application of 1.0 lb. soluble N/1000 ft^2 would result in very high total N rates. For example, a natural organic fertilizer with 10% soluble N and 90% slow release N would yield a total N application rate of 10 lbs. N/1000 ft^2 when applied at 1.0 lb of soluble N per 1000 ft^2 , which far exceeds recommended annual rates of N for typical turfgrasses. Thus, for slow release fertilizers with little soluble N, the maximum amount applied in one application should not exceed total annual N requirements of the turf.

As previously discussed, the annual turfgrass requirements for N vary considerably depending on a variety of conditions. Most critical, however, is turfgrass species. The annual N requirements for maintaining established stands of the most common turfgrass species grown in Maryland generally fall into the ranges listed in Table 1.

Table 1. Nitrogen Rate Recommendations for Commercially Maintained Turfgrass

Total Nitrogen Annually (lbs. N/1000 ft²)		
Cool Season Grasses	Years 1-2	Subsequent Years
Kentucky bluegrass	3.0 – 4.5	3.0 – 4.0
Turf-type tall fescue	3.0 – 4.0	2.0 – 3.0
Fine fescue	1.0 – 3.0	0 – 2.0
Perennial Ryegrass	3.0 – 4.0	3.0 – 4.0
Warm Season Grasses	Years 1-2	Subsequent Years
Bermudagrass	3.0 – 4.0	3.0 – 4.0
Zoysiagrass	1.0 – 3.0	0 – 2.0

Numerous factors influence whether moderate adjustments to these rates may be warranted. For example, if clippings are returned to the site when it is mowed, reductions in the annual N rates (as well as P and K) may be possible. Also, if the site receives little use, and thus does not need higher growth rates to recover from traffic, lower rates may be advisable. Other means of possible reductions in total N requirements include the use of iron, increasing the height of mowing, and careful selection of cultivars when seeding, overseeding or sodding.

Conversely, several factors may warrant moderate increases in annual rates. These include heavily used sites that need high recuperative rates, sites that are mowed lower than recommended due to use requirements, and sites that have been damaged from adverse environmental conditions or pests. Also, on sites where pesticide use is not economically feasible or permitted, somewhat higher N rates can be important in minimizing many pest problems, particularly weeds and diseases. Rates much higher than recommended, however, can have the opposite effect.

Timing of Nitrogen Applications

The primary potential for N loss from turfgrass sites is when excessive rates of N, particularly NO₃-N, are applied to turf that is not actively growing. Thus, most of the annual fertilizer requirement should be applied during periods of active shoot (leaf blades, rhizomes, stolons) and/or root growth (Table 2).

Table 2. Recommended Periods for N Fertilization of Commercially Maintained Turf

	Recommended Periods	Periods to Avoid
Warm Season Grasses	1 month before dormancy breaks (mid-April in central MD; later in western MD and earlier on the eastern shore) through September 1 st	September 1 st through 1 month before dormancy breaks During severe or prolonged drought
Cool Season Grasses	1 month before topgrowth starts (late March in central MD) through early June Late August through 6 weeks after first killing frost	Mid-June through mid-August When turf is dormant due to heat, drought, or cold

The primary period for growth of warm season grass species (zoysiagrass, bermudagrass, and buffalograss) is from mid-spring, after dormancy has broken, through mid-fall, when the first killing frost is experienced. Thus, N applications to warm season grasses should generally be restricted to these periods. However, fertilizer that contains primarily NH₄-N can be applied up to a month before dormancy is typically broken in the spring so that N is available for plant uptake at this time. Applications after September 1 are not generally recommended due to the possible enhancement of winterkill, particularly with bermudagrass. However, if bermudagrass has been overseeded with a cool season species such as perennial ryegrass, up to 1.0 lb. N/1000 ft² may be applied after September 1 to enhance its performance.

Cool season grasses have a longer growth period. They can exhibit growth at virtually any time during the year if moisture and temperature conditions are conducive. The prime periods for growth are typically from late winter through early summer and from late summer through early winter. Research would indicate that 2/3 to 3/4 of the total annual N should be applied during the latter period to maximize cool season turfgrass performance and quality.

Under extended hot and dry periods during mid-summer, cool season grasses may experience a period of dormancy until rainfall occurs. Nitrogen fertilizer should not be applied at this time. If irrigation is available or if rainfall is adequate throughout the summer, little dormancy will occur and N uptake of cool season grasses will continue. Although not generally needed, applications of ¼ to ½ lb./1000 ft² can be made to these sites if growth is not adequate to meet the demands of the use of the site.

During the winter months, although top growth may have virtually ceased, root growth and N uptake may still occur during the periods when the ground is not actually frozen, particularly with Kentucky bluegrass. Research has shown that applications of N during this period can enhance root growth and spring performance of turf. Also, some sites tend to be excessively wet each spring, making fertilization with large equipment difficult. Fertilization earlier when the ground is firmer may be the best alternative.

There is minimal risk of runoff or leaching problems from winter application of N if the following guidelines are followed: No more than ½ - 1.0 lbs. N/1000 ft² should be applied, fertilizers containing significant amounts of NO₃-N should not be used, and applications should not be made to frozen ground if significant rainfall is in the immediate forecast. For most situations, however, mid-winter applications are not necessary and the guidelines listed in Table 2 should be followed.

Phosphorus and Potassium

Phosphorus (P) is critical in the establishment of turfgrass. Inadequate soil P will result in very poor seedling vigor, slow establishment of grass, and a stand with very poor density and root growth. Thus, soil will be much more susceptible to erosion. Weed encroachment will also be much more severe due to the lack of competition from the thin turfgrass stand. Thus, it is essential that sufficient P be added to the soil at the time of seeding if soil levels are inadequate. Although not as critical as during establishment of turfgrass, deficiencies of P in mature turf can result in poor spring greenup, reduced vigor, reduced density, and reduced drought tolerance. Light applications of P are generally sufficient to overcome deficiencies in mature turf.

Potassium (K) is not as critical as N or P during the initial establishment phase of turf. However, K plays an important role in mature turf regarding rhizome production and tolerances to heat, drought, and cold. Thus, sufficient K needs to be available for turfgrass to ensure that quality turf will be obtained during and after summer and/or winter stresses. Severe deficiencies of K will result in thin, chlorotic turf that may also exhibit a lack of vigor in turf.

Whereas N application recommendations cannot currently be obtained from soil tests, recommendations for application of P and K should be obtained from such tests. A commercial turf maintenance company should test sites within a year of the initiation of management of the site. Until such time as the first soil test is taken, not more than 1.0 lb. P₂O₅ and 2.0 lb. K₂O per 1000 ft² should be applied for maintenance of turf. After the initial soil test, subsequent sampling every 3 to 4 years is generally sufficient to monitor soil P and K levels.

Sites having different soil types, sites with different use or management histories, and/or sites having substantially different fertility levels as determined by previous soil tests should be sampled separately. Conversely, sites having similar soil types, having similar use and management histories, and having similar fertility levels as determined by past soil tests may be lumped together into one sample. For example, a single sample may be sufficient for an athletic field complex (or town house development, etc.) that has had similar management over a period of time and has a relatively uniform soil type.

Table 3. P and K Recommendations for Commercially Maintained Turf (Based on U.M. Soil Test Results)

	Soil Test Category		
	low	medium	optimum----excessive
	lbs. P ₂ O ₅ or K ₂ O/1000 ft ²		
Phosphorus	2.0	1.0	0
Potassium	2.0 – 4.0	1.0 - 2.0	0 – 1.0

Soil Reaction

Maintaining soil pH in an optimum range is important for maximizing the efficiency of nutrient use, and can be important in reducing weed and disease problems. Turfgrass can withstand a rather broad range of soil pH, but 5.8 to 6.4 is generally considered ideal. Wide deviations from this range can result in reduced P and micronutrient availability, and can interfere with soil N metabolism and availability. Depending on turfgrass species, problems in turf may start to occur at soil pH above 7.8 and below 5.4. Thus, to maximize efficiency of nutrient availability and use, soil tests should be taken as recommended previously for soil P and K to determine soil pH.

Recommended limestone applications to achieve a soil pH of about 6.4 are shown in Table 4. If diseases such as take-all patch of bentgrass, summer patch of Kentucky bluegrass, or spring dead spot of bermudagrass are of concern, maintaining lower soil pH (5.4 - 5.7) may be desirable, and reduced or no limestone should be applied to achieve this level. Also, it is recommended, when practical, that limestone be applied approximately one month or more before seeding to minimize potential P availability problems and the potential for volatilization loss of applied N.

It should be emphasized that the information presented within this publication for N, P, K, and limestone applications is meant only as a guideline. While these recommendations should result in satisfactory sod production in most situations, there are many factors that could impact whether modifications of these recommendations are warranted for a specific site. Higher rates of fertilization may be warranted for specific sites and/or conditions.

Table 4: Limestone Maintenance Recommendations (Pounds per Acre)

soil pH			Coastal	Plain	Piedmont &	Mountain
	loamy sands	sandy loams	loams	silt loams and silty clay loams	loams	silt loams and silty clay loams
6.4	0	0	0	0	0	0
6.3	0	0	0	0	545	650
6.2	0	0	545	545	760	1090
6.1	0	545	650	760	980	1415
6.0	435	760	870	980	1195	1850
5.9	545	870	1090	1195	1415	2180
5.8	650	1090	1195	1415	1740	2505
5.7	760	1195	1415	1630	1960	2940
5.6	870	1305	1630	1850	2180	3265
5.5	980	1525	1850	2070	2395	3590
5.4	1090	1630	2070	2180	2720	3920
5.3	1195	1850	2180	2395	2940	4355
5.2	1305	1960	2285	2610	3155	4355
5.1	1415	2070	2505	2830	3375	4355
5.0	1525	2285	2720	3050	3590	4355
4.9	1630	2395	2940	3155	3810	4355
4.8	1740	2505	3050	3375	4135	4355
4.7	1850	2720	3265	3590	4355	4355
4.6	1960	2830	3375	3810	4355	4355
4.5	2070	3050	3590	4030	4355	4355

- 1) divide the above rates by 43.5 to obtain the equivalent rates in pounds per 1000 square feet
- 2) do not apply more than 2,000 pounds per acre for any one maintenance application to turfgrass

Related Publications: TT-114 Nutrient Management Guidelines for Sod Production in Maryland

Educating People to Help Themselves Local Governments • U.S. Department of Agriculture Cooperating

The University of Maryland is equal opportunity. The University's policies, programs and activities are in conformance with pertinent Federal and State laws and regulations on nondiscrimination regarding race, color, religion, age, national origin, sex, and disability. Inquiries regarding compliance with Title VI of the Civil Rights Act of 1964, as amended; Title IX of the Educational Amendments; Section 504 of the Rehabilitation Act of 1973; and the Americans With Disabilities Act of 1990; or related legal requirements should be directed to the Director of Personnel/Human Relations, Office of the Dean, College of Agriculture and Natural Resources, Symons Hall, College Park, MD 20742.